## **Post-traumatic Stress Disorder**

Ros, T. et al. Neurofeedback Tunes Scale Free Dynamics in spontaneous Brain Activity. Cerebral Cortex. 2016 Sep 12.

Nicholoson, A.A. et al. Alpha oscillation neurofeedback modulates amygdala complex connectivity and arousal in posttraumatic stress disorder. Neuroimage Clin. 2016 Jul 14;12,506-516

Tursich, M. et al. Distinct intrinsic network connectivity patterns of post-traumatic stress disorder symptom clusters. Acta Psychiatr Scand. 2015 Jul;132(1)29-38

Kluetsch.R.C. et al. Plastic modulation of PTSD resting-state networks and subjective wellbeing by EEG neurofeedback. Acta Psychiatr Scand. 2014 Aug; 130(2),123-136

Van der Kolk, B.A. et al. A randomized controlled study of neurofeedback for chronic PTSD. <u>PLOS ONE</u> 2016, Dec 1-18.

Clancy, K. et al. Restless 'rest': intrinsic sensory hyperactivity and disinhibition in post-traumatic stress disorder. Brain 2017, June 3, doi 10.1093.

Olson, E.A. et al. Disruption of white matter structural integrity and connectivity in posttraumatic stress disorder: A TBSS and tractography study. Depress Anxiety 2017 May; 34(5): 437-445.

Falconer, E. et al. The neural networks of inhibitory control in posttraumatic stress disorder. <u>J Psychiatry Neurosci</u> 2008 Sep; 33(5): 413-422.

Clancy, K., et al., Restless Rest: Intrinsic Sensory Hyperactivity and Disinhibition in Post Traumatic Stress Disorder. Brain 2017, 1-10

National Institute of Mental Health. Brain Biomarkers Could Help Identify those at Risk of Severe PTSD. Science Daily January 28, 2019

Van der Kolk, B., et al. A Randomized Controlled Study of Neurofeedback for Chronic PTSD. <u>PLOS ONE</u> December 16, 2016

Wang, C. et al. Utility of P300 ERP in monitoring post trauma mental health: a longitudinal study in military personnel returning from combat deployment. Journal of Psychiatric Research 2018, 101, 5-13